



SUMMER SCHOLARSHIP - 2005/2006 SEASON

Project title: Genetic variation in cotton for tolerance to waterlogged conditions

Aims and milestones:

- Determine the leaf nutrient level, leaf colour, leaf photosynthesis, plant and root morphology response of cotton under waterlogged conditions
- Evaluate the physiological responses of fourteen cotton cultivars under waterlogged conditions.
- Evaluate methodologies to measure the response of cotton to waterlogged conditions
- Build on previous findings to improve the understanding of waterlogging effects on cotton physiology in terms of tissue structure and function, and plant growth and development.

Staff:

Project supervisors: Dr Greg Constable; Prof Bruce Sutton; Dr Daniel Tan

Position in organisation: Program Leader (CSIRO Plant Industry, Cotton Research Unit); Chair of the Academic Board, University of Sydney; Lecturer in Agronomy (Farming Systems), Faculty of Agriculture, Food and Natural Resources, University of Sydney

Telephone (02) 6799 1522; (02) 9351 3183; (02) 9036 9014

Facsimile (02) 6793-1186; (02) 9351 3572; (02) 9351 4172

Postal Address Locked Bag 59 Narrabri NSW 2390; A14 – Quadrangle, The University of Sydney, NSW 2006 Australia; A20 - John Woolley, The University of Sydney, NSW 2006 Australia

Project Summary

This summer scholarship project was undertaken by Warren Conaty. Warren is in his fourth and final year of Bachelor of Science in Agriculture at the University of Sydney and is majoring in Agronomy. The scholarship commenced on 10th January, 2006 and was completed on 22nd February 2006.

Background

Waterlogging is a world-wide phenomenon that affects both the distribution of plants in their natural ecosystems as well as crop yield in agricultural regions. Waterlogging can severely affect cotton growing in the heavy clays (Vertosols) in Australia. The annual production losses due to waterlogging in Australia is \$A180 million (Price 1993). Cotton production in Australia is:

- focused in regions with heavy clay soils with inherent low drainage rates;
- characterised by field with low gradients;
- almost exclusively furrow irrigated; and

As cotton is grown in summer dominant rainfall regions, which have a high probability of intense summer storms, there is a high likelihood of disruption of irrigation schedules and occurrence of waterlogged conditions. Waterlogging can result in yield reductions of up to 10% (Bange *et al.* 2004).

Waterlogging occurs in saturated soils, when the air filled porosity (AFP) falls below 10% (Hodgson 1982). In a well drained soil, the air filled porosity usually ranges from 10 to 40% of the total soil volume. However, waterlogging reduces these gas-filled pores, substantially diminishing the supply of oxygen to the roots resulting in hypoxia. This occurs due to the prevention of the replacement of oxygen consumed in the soil.

Waterlogging has a wide range of effects on cotton including decreased root and shoot biomass as well as an increase in root and leaf senescence (Bange *et al.* 2004; Hodgson and Chan 1982). Waterlogging also affects nutrient availability, uptake, translocation and leaf nutrient concentration (Hocking *et al.* 1985). Furthermore, hypoxia results in decreased stomatal conductance and leaf water potential, and hence, reduced photosynthesis.

Aims and objectives

This study evaluates the physiological responses of fourteen cotton genotypes under waterlogged and non-waterlogged irrigated conditions in relation to leaf nutrient level, leaf colour, leaf photosynthesis, plant morphology and final yield.

Methodology

Thirteen different upland cotton (*Gossypium hirsutum*) and one pima cotton genotype (*Gossypium barbadense*) from a range of natural environments were subjected to waterlogged and non-waterlogged irrigated (control) conditions at the ACRI, Narrabri (see Table 1). Control plots were irrigated based on Best Management Practices. Waterlogged plots were subjected to optimal irrigation schedules plus 48 to 72 hours of additional irrigation. The physiological responses of these cotton genotypes were assessed in relation to leaf nutrient status, leaf colour, leaf photosynthetic rate, plant morphology and crop yield. More regular measurements were undertaken on four cotton genotypes (Georgia King, LA 887, Sicot 71 and Sicot 73) with variable responses to waterlogged conditions.

Table 1. Cotton genotypes, origin and environmental conditions used in this study

Variety	Origin	Environment	Soil Type
CIM 443	Pakistan	Dry	Clay
Coker 315	USA- Texas	Cool	Clay
Gohar 87	India	Dry	Clay
Sicot 71	Australia- CSIRO	Dry	Clay
Sicot 73	Australia- CSIRO	Dry	Clay
Sicot 80	Australia- CSIRO	Dry	Clay
Codetec 401	Brazil	Wet	Loam
Deltapine 16	USA- Mississippi	Humid	Silt
Deltapine 90	USA- Arizona	Hot	Loam
Georgia King	USA- Georgia	Humid	Silt
LA 887	USA- Louisiana	Humid	Silt
McNair 1032	USA- South Carolina	Humid	Silt
PD93057	USA- South Carolina	Humid	Silt
Pima A-8	USA- Arizona	Hot	Loam

Data Collection

Leaf Nutrient Status

The first fully expanded leaf was collected at regular intervals, dried and analysed via Inductively Coupled Plasma Mass Spectrometry (ICP-MS) for phosphorus (P), potassium (K), iron (Fe), manganese (Mn), magnesium (Mg), calcium (Ca), copper (Cu), zinc (Zn), boron (B), aluminium (Al) and sodium (Na). Leaf colour was measured using the Minolta Spad-502. Leaf photosynthetic rates were measured 24 hours before waterlogging and then again 24 hours after the irrigation event for the control and during waterlogging. Boll retention and yield (kg seed/ha) were measured.

Results

Leaf Nutrient Status

Leaf blade concentrations of phosphorus varied between genotypes ($P < 0.001$). Phosphorus concentrations decreased due to waterlogging except for the genotypes Sicot 80, PD93057, Sicot 73 and Sicot 71.




Figure 1. Phosphorus concentrations (mg/kg) in leaves of irrigated and waterlogged genotypes.

Leaf blade concentrations of potassium varied between genotypes ($P < 0.001$). Potassium concentrations decreased due to waterlogging except for Coker315, Pima A-8, PD93057, Codetec 401, McNair 1032, Sicot 71, Gohar 87 and Sicot 73.




Figure 2. Potassium concentrations (mg/kg) in leaves of irrigated and waterlogged genotypes.

Leaf iron concentration increased in all genotypes due to waterlogging (Figure 3). Genotypes displayed different concentrations of iron ($P < 0.001$). Hence, leaf chlorosis (yellowing) in the waterlogged treatments (Figure 6), was not due to low leaf iron concentrations. This anomaly could be due to waterlogged conditions that reduce the available forms of iron (Fe^{2+}) and increase unavailable iron (Fe^{3+} and others) concentrations.




Figure 3. Iron concentrations (mg/kg) in leaves of irrigated and waterlogged genotypes.

Photosynthetic Rate

Pre-irrigation, photosynthetic rates were higher in waterlogged treatments ($P < 0.01$) due to higher soil moisture availability (data not shown) (Figure 4). However, similar effects were observed between the cultivars ($P = 0.652$) in the same treatment.




Figure 4. Leaf photosynthetic rates of genotypes LA 887 and Sicot 71 before irrigation.

Following saturation, photosynthetic rate was higher ($P < 0.05$) in the LA 887 control compared to the waterlogging treatment (Figure 5). However, photosynthetic rates for Sicot 71 were not different across irrigation treatments ($P = 0.693$).




Figure 5. Leaf photosynthetic rates of the genotypes LA 887 and Sicot 71 post irrigation (control) and during waterlogging.

Leaf Colour

Leaf colour was strongly affected by waterlogging, resulting in yellowing of leaves ($P < 0.001$, Figure 6). However some genotypes were not as affected by waterlogging. The genotypes that had no difference in leaf colour response to waterlogging were Sicot 71, Sicot 73, Coker 315, Sicot 80 and Pima A-8.




Figure 6. Relative leaf colour for waterlogged and control irrigated genotypes.

Boll Retention and Yield

There was a higher ($P < 0.05$) boll retention rate in the control compared to the waterlogging treatment in all cultivars except Sicot 71 (Figure 7). Yield results (kg seed/ha) are shown in Figure 8. Waterlogging significantly reduced the yield (kg/ha seed) of waterlogging sensitive varieties LA 887 and Georgia King.




Figure 7. Boll retention rates for Georgia King, LA 887, Sicot 71 and Sicot 73 under waterlogged and control irrigated conditions.




Figure 8. Yield (kg seed/ha) for waterlogged and control irrigated genotypes (Note: 1kg seed/ha is approximately equal to 40kg ginned lint/ha).

Discussion

Leaf blade nutrient concentration

Waterlogging reduced both phosphorus and potassium leaf blade concentrations. This can be attributed to inhibited nutrient uptake under anaerobic conditions (Trought and Drew 1980). These deficiencies in waterlogged plants also occur due to the inhibition of ion uptake and the translocation of nutrients from older leaves to supply the nutrients to the younger leaves (Drew 1983). At present, there have been no studies that investigate whether waterlogging impacts on the population of active arbuscular mycorrhizal fungi (AMF). Such a study is necessary in order to determine if waterlogging adversely affects AMF populations, and thus, the potential phosphorus uptake.

Waterlogging increased the iron concentration in all genotypes. Numerous studies on cotton and other field crops have shown an increase in iron concentration following waterlogging (Hodgson 1990). This increase may be due to the effect of increased iron availability in the soil. When a soil is inundated, ferric iron (Fe^{3+}) is reduced to the more soluble ferrous (Fe^{2+}) ions, increasing soil iron availability. This chemical reaction is characterised by the transformation of the brown colour in an iron-rich soil to shades of grey as large amounts of iron ions are brought into solution (Ponnamperuma 1984). This effectively increases the availability of these elements, which can reflect an increase in total iron concentration of waterlogged plants.

However, acute deficiencies of iron can still occur in waterlogged cotton. This is because iron nutrition should reflect the active chemical form of iron (Fe^{2+}), rather than total iron content. Therefore total iron concentrations can be misleading (Figure 3) as it is the active form of iron, Fe^{2+} that is critical for plant nutrition and chlorophyll production (Hodgson 1990). The inherent problem in iron nutrition is that the determination of the Fe^{2+} content cannot be achieved by a commercial laboratory, and requires the use of fresh leaves, analysed a few hours after sampling.

An iron deficient plant will show signs of interveinal yellowing. The veins usually remain green, but if the deficiency is extreme, the leaf can become totally chlorotic and turn white. Ironically, the total iron concentration in chlorotic leaves is often similar to, or higher than in green leaves. This is because the majority of this iron is unavailable for use by plants as it is in the inactive Fe^{3+} form (Hodgson 1990). Iron deficiencies occur in cotton because as the soil becomes increasingly waterlogged, the diffusion of CO_2 gas out of the soil is reduced,

creating a build up of carbon dioxide (CO₂) in the soil. This results in the formation of bicarbonate ions, increasing both the soil pH and the alkalinity of the leaf tissue. Under these conditions, available ferrous iron (Fe²⁺) is reduced to inactive forms of iron (predominantly ferric or Fe³⁺) (Hodgson 1990). For each unit increase in leaf pH, the availability of iron is decreased approximately 100-fold (Hodgson 1990). Therefore, waterlogging results in the reduction of available iron which the plant must convert into ferrous iron (Fe²⁺) before it can be used. This process is metabolically costly to a plant already suffering from reduced metabolism associated with hypoxia, resulting in iron deficiencies.

Foliar application of iron (II) sulphate on cotton can help ameliorate the effects of iron chlorosis, returning cotton foliage to its normal colour. In addition, foliar sprays of iron chelates applied to crops before irrigation and subsequent waterlogging events can help prevent iron deficiencies. Despite the risk of foliar burn, a single spray on cotton at Narrabri increased lint yields by 6%, and two foliar sprays increased lint yield by 11% (Hodgson 1990).

Photosynthetic Rate and Leaf Colour

Photosynthetic rates in response to flooding were reduced by 19% in the susceptible genotype (LA 887) and 7% in the more tolerant genotype (Sicot 71). Similar results were observed by Meyer *et al.* (1987) who reported that photosynthetic rates in cotton decreased by 16% following seven days of waterlogging. Flooding and subsequent waterlogging of the soil is usually followed by a rapid reduction in the rate of photosynthesis in waterlogging susceptible plants (Kozłowski and Pallardy 1984). Photosynthesis of waterlogged plants is reduced initially by stomatal closure and later by a reduction in photosynthetic capacity (Kozłowski and Pallardy 1984). Plants under waterlogging stress have reduced stomatal aperture, and transpiration rates. This helps prevent waterlogging-induced wilting and thus reduced photosynthetic rate.

Chlorosis of leaves, resulting from a reduction in the level of chlorophyll in the leaf, is often one of the early symptoms of waterlogging. Leaf yellowing to some extent was observed in all genotypes in the waterlogging treatment. As chlorophyll is the light capturing pigment in plants, a reduction in its level in the leaf will reduce photosynthetic rate. Decreased chlorophyll content due to waterlogging has been reported in numerous *Brassica* species (Ashraf and Mehmood 1990), barley (Pang *et al.* 2004), maize (Bragina *et al.* 2004), field pea (Ladygin 2004) and okra (*Hibiscus esculentus*) (Ashraf and Arfan 2005). The inhibition of chlorophyll production in leaves is at least partly responsible for a reduction in photosynthetic rates (Bragina *et al.* 2004). The importance of chlorophyll content to photosynthesis is emphasised by a parallel decline in both chlorophyll and photosynthetic rate (Kozłowski and Pallardy 1984).

Boll Retention and Yield

Waterlogged conditions affect almost every aspect of shoot behaviour and thus, final crop yield. The effects of waterlogging on the shoot are due to modifications in internal flow of water, hormones, toxins, inorganic nutrients and photosynthates from the root to the shoot. Waterlogging reduced boll retention in the genotypes Georgia King by 23%, LA 887 by 24% and Sicot 73 by 24%. No difference in boll retention was observed in Sicot 71 as a result of waterlogging. These results are similar to those observed by Bange *et al.* (2004). According to Bange *et al.* (2004) dry matter accumulation was reduced by 32% due to a 35% reduction in radiation use efficiency. The reduction in dry matter (32%) resulted in decreased yield, achieved principally through decreased boll retention, and hence, reduced boll number. These reductions in boll retention rates due to waterlogging reduced crop yields. Waterlogging significantly reduced the yield (kg/ha seed) of waterlogging sensitive varieties LA 887 by 21%, and Georgia King by 17%. In contrast, yields of genotypes that were more tolerant to waterlogged conditions were only reduced by 4% for Sicot 71 and 3% for Sicot 80. These differences in yield were closely related to boll retention rates.

Conclusion

Waterlogging reduced vegetative growth, dry matter accumulation and subsequently, crop yield. The physiological consequences of waterlogged conditions include reduced photosynthetic rate, reduced water and nutrient uptake, and altered shoot and root hormonal status.

There was some genetic tolerance to waterlogged conditions when comparing the physiological response of the more susceptible varieties (LA 887 and Georgia King) with local varieties (Sicot 71, Sicot 73 and Sicot 80) which appeared to be more tolerant. Thus direct selection for yield in the local breeding program has appeared to have indirectly selected for increased tolerance to waterlogging as new local varieties had only 3% to 4% reduced yield with waterlogging treatments in this experiment. Until this study, there have been few studies investigating the genetic variation in the response of cotton genotypes to waterlogging stress. The knowledge obtained from further studies into these areas may be utilised in agronomic field management practices and conventional and transgenic breeding programs. Manipulation of genes controlling these morphological adaptations has the potential to enable sensitive plant species to endure longer periods of waterlogging before cellular functions cease due to a lack of energy.

However, it is important that correct agronomic management and best management practices are still observed in relation to waterlogging. Improvements to field gradients through laser levelling, shortening of furrow lengths and the implementation of two metre beds are some methods to reduce the extent of waterlogging stress in furrow irrigation. Furrow irrigation efficiency can be increased through the use of larger siphons and the broadening of tail drains. This will reduce the inundation period and therefore, the potential period of waterlogging stress.

Presentations and public relations

A media release was compiled and gave a brief overview of the summer scholarships being undertaken for the summer of 2005/2006 and the students associated with each scholarship (Cotton Magazine, The Courier – 17 Jan 06).

A presentation was given to staff of the Australian Cotton Research Institute, on 22nd February 2006 following completion of the project.

A presentation was given to the board of the Cotton (CCC) CRC at Darlington Board Room, The University of Sydney on 27 July 2006.

An article “Genetic variation in cotton for tolerance to waterlogged conditions” was published in the ACGRA 13th Cotton Conference held in the Gold Coast from 8-10 August 2006.

References

Ashraf M, Arfan M (2005) Gas exchange characteristics and water relations in two cultivars of *Hibiscus esculentus* under waterlogging. *Biologia Plantarum* **49**, 459-462.

Ashraf M, Mehmood S (1990) Effects of waterlogging on growth and some physiological parameters of four *Brassica* species. *Plant and Soil* **121**, 203-209.

Bange MP, Milroy SP, Thongbai P (2004) Growth and yield of cotton in response to waterlogging. *Field Crops Research* **88**, 129-142.

Bragina TV, Ponomareva YV, Drozdova IS, Grinieva GM (2004) Photosynthesis and dark respiration in leaves of different ages of partly flooded maize seedlings. *Russian Journal of Plant Physiology* **51**, 342-347.

Drew MC (1983) Plant injury and adaptation to oxygen deficiency in the root environment: a review. *Plant and Soil* **75**, 179-199.

Hocking PJ, Reicosky DC, Meyer WS (1985) Nitrogen status of cotton subjected to two short term periods of waterlogging of varying severity using a sloping plot water table facility. *Plant and Soil* **87**, 375-391.

Hodgson AS (1982) The effects of duration, timing and chemical amelioration of short-term waterlogging during furrow irrigation of cotton in a cracking grey clay. *Australian Journal of Agricultural Research* **33**, 1019-1028.

Hodgson AS (1990) Micronutrients: Are they important under waterlogging? In 'Fifth Australian Cotton Conference'. Broadbeach pp. 165-170. (ACGRA).

Hodgson AS, Chan KY (1982) The effect of short term waterlogging during furrow irrigation of cotton in a cracking grey clay. *Australian Journal of Agricultural Research* **33**, 109-116.

Kozlowski TT, Pallardy SG (1984) Effect of flooding on water, carbohydrate and mineral relations. In 'Flooding and plant growth'. (Ed. TT Kozlowski) pp. 165-193. (Academic Press: London).

Ladygin VG (2004) The effect of root hypoxia and iron deficiency on the photosynthesis, biochemical composition, and structure of pea chloroplasts. *Russian Journal of Plant Physiology* **51**, 28-40.

Meyer WS, Reicosky DC, Barrs HD, Smith RCG (1987) Physiological responses of cotton to a single waterlogging at high and low N-levels. *Plant and Soil* **102**, 161-170.

Pang JY, Zhou MX, Mendham N, Shabala S (2004) Growth and physiological responses of six barley genotypes to waterlogging and subsequent recovery. *Australian Journal of Agricultural Research* **55**, 895-906.

Ponnamperuma FN (1984) Effects of flooding on soils. In 'Flooding and plant growth'. (Ed. TT Kozlowski) pp. 10-45. (Academic Press: London).

Price P (1993) Resource base: the nation's vital asset. *Agricultural Science* **6**, 42-45.

Trought MCT, Drew MC (1980) The development of waterlogging damage in wheat seedlings (*Triticum aestivum* L.) II. Accumulation and redistribution of nutrients by the shoot *Plant and Soil* **56** 187-199.